

Decomposition of vector-valued divergence free Sobolev functions and shape optimization for stationary Navier-Stokes equations *

Gengsheng Wang

School of Mathematics, Wuhan University,
Wuhan, Hubei, 430072, P.R.of China
wanggs@public.wh.hb.cn

Donghui Yang

School of Mathematics, Wuhan University,
Wuhan, Hubei, 430072, P.R.of China
dongfyang@yahoo.com.cn

Abstract

We establish a divergence free partition for vector-valued Sobolev functions with free divergence in $\mathbf{R}^n$, $n \geq 1$. We prove that for any domain Ω of class $\mathcal{C}$ in $\mathbf{R}^n$, $n = 2, 3$, the space $D_0^1(\Omega) \equiv \{\mathbf{v} \in H_0^1(\Omega)^n; \operatorname{div} \mathbf{v} = 0\}$ and the space $H_{0,\sigma}^1(\Omega) \equiv \overline{\{\mathbf{v} \in C_0^\infty(\Omega)^n; \operatorname{div} \mathbf{v} = 0\}}^{\|\cdot\|_{H^1(\Omega)^n}}$, which is the completion of $\{\mathbf{v} \in C_0^\infty(\Omega)^n; \operatorname{div} \mathbf{v} = 0\}$ in the $H^1(\Omega)^n$ -norm, are identical. We will also prove that $H_{0,\sigma}^1(D \setminus \overline{\Omega}) = \{\mathbf{v} \in H_{0,\sigma}^1(D); \mathbf{v} = 0 \text{ a.e. in } \Omega\}$, where D is a bounded Lipschitz domain such that $\Omega \subset \subset D$. These results, together with properties for domains of class $\mathcal{C}$, are used to solve an existence problem in the shape optimization theory of the stationary Navier-Stokes equations.

Key words. Partition of functions, shape optimization, stationary Navier-Stokes equation

AMS subject classifications. 35Q30, 49Q30, 49J10.

1. Introduction

*This work was supported by National Natural Science Foundation of China, Grant No. 10471053, 60574017 and by the key programme of the Ministry of National Education of China.

In this paper, we study an existence problem in the shape optimization theory of the stationary Navier-Stokes equation over $D \setminus \overline{\Omega}$ with $\Omega \subset\subset D$ varying. Here, $D \subset \mathbf{R}^n$, $n = 2, 3$, is a given bounded Lipschitz domain and Ω is supposed to satisfy the property $\mathcal{C}$ to be defined in section 2. Our main ingredient for such an investigation is a new divergence-free partition for divergence free vector-valued Sobolev functions defined over $D \setminus \overline{\Omega}$. This partition is established by making use of the classical Hodge Theory in Differential Geometry. To be more precise, we let D be an open subset in $\mathbf{R}^n$ ($n \geq 1$), and let $\{U_j\}_{j=1}^m$ be an open covering of $\overline{D}$. Suppose that $\mathbf{u} \in D_0^1(D) \equiv \{\mathbf{u} \in H_0^1(D)^n; \operatorname{div} \mathbf{u} = 0\}$. Then our first main result states that there are functions $\{\mathbf{u}_j\}_{j=1}^m$ such that $\mathbf{u}_j \in D_0^1(U_j)$, $\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j$ over D and $\|\mathbf{u}_j\|_{H^1(U_j)^n} \leq C \|\mathbf{u}\|_{H^1(D)^n}$ for some positive constant C independent of $\mathbf{u}$. Moreover, such a partition inherits the following local property: If $\mathbf{u}|_{\Omega} = 0$ with $\Omega \subset\subset D$ and Ω is of class $\mathcal{C}$ to be defined in section 2, then we can choose $\{\mathbf{u}_j\}_{j=1}^m$ such that $\mathbf{u}_j|_{\Omega} = 0$ for $j = 1, \dots, m$.

The above decomposition, besides its application to the shape optimization problem for the stationary Navier-Stokes equation, has many corollaries which might be interesting in their own right. For instance, it gives the following property for the function spaces $H_{0,\sigma}^1(\Omega)$ and $H_{0,\sigma}^1(D \setminus \overline{\Omega})$ where

$$H_{0,\sigma}^1(\Omega) \equiv \overline{\{\mathbf{u} \in C_0^\infty(\Omega)^n; \operatorname{div} \mathbf{u} = 0\}}^{\|\cdot\|_{H^1(\Omega)^n}}$$

(the completion of $\{\mathbf{u} \in C_0^\infty(\Omega)^n; \operatorname{div} \mathbf{u} = 0\}$ in the norm of $H^1(\Omega)^n$):

If Ω is of class $\mathcal{C}$ and D is Lipschitz, then

$$H_{0,\sigma}^1(\Omega) = D_0^1(\Omega), \quad (1.1)$$

$$H_{0,\sigma}^1(D \setminus \overline{\Omega}) = \{\mathbf{v} \in H_{0,\sigma}^1(D); \mathbf{v} = 0 \text{ a.e. in } \Omega\}. \quad (1.2)$$

(1.2) has the following consequences:

(i) If $\mathbf{v} \in H_{0,\sigma}^1(D \setminus \overline{\Omega})$, then $\tilde{\mathbf{v}} \in H_{0,\sigma}^1(D)$, where

$$\tilde{\mathbf{v}} = \begin{cases} \mathbf{v} & \text{in } D \setminus \overline{\Omega}, \\ 0, & \text{in } \Omega. \end{cases}$$

(ii) If $\mathbf{v} \in H_{0,\sigma}^1(D)$ and $\mathbf{v} = 0$ a.e. in Ω , then $\hat{\mathbf{v}} \in H_{0,\sigma}^1(D \setminus \overline{\Omega})$, where $\hat{\mathbf{v}}$ is the restriction of $\mathbf{v}$ over $D \setminus \overline{\Omega}$.

(1.1) and (1.2) can be compared with the classical result in [7]: If $v \in H_0^1(D)$ and $v = 0$ quasi-everywhere in $D \setminus \Omega$ (where Ω is an open set), then $v \in H_0^1(\Omega)$. They can also be compared with the result in [14], which states that if $v \in H_0^1(D)$ and $v = 0$ a.e. in $D \setminus \Omega$ (where Ω is of class $\mathcal{C}$), then

$v \in H_0^1(\Omega)$. We notice that (1.1)-(1.2) have been obtained earlier in [6] and [11] under the assumption that Ω is Lipschitz. The novelty in (1.1)-(1.2) is that Ω does not have to have a Lipschitz boundary. In fact, a generic domain with property $\mathcal{C}$ is not a finite union of Lipschitz domains. It may have cusps and infinitely many oscillations with vanishing amplitude.

Fundamentally based on the aforementioned decomposition and properties on the family $\mathcal{O}$ which will be defined in Section 2, we study in section 5 a shape optimization problem. In this problem, we will look for a domain of class $\mathcal{C}$ which minimizes a certain cost functional associated with the solutions of the stationary Navier-Stokes equation. Our main purpose for this part of the paper is to prove Theorem 5.1 to be stated in section 5.

There have been extensive studies for the existence problems in the shape optimization theory for elliptic equations in the past twenty years. Here, we mention the work in [4], [10], where results are obtained over domains with sufficiently smooth boundary. We mention the work in [12] and [3], which is based on a certain capacity constraint and penalty terms, respectively. For more recent work, we mention [14] and [15] where the domains are assumed to have a certain segment property. These two papers are based on the analysis of the set of bounded open sets of class $\mathcal{C}$ in the sense of Maz'ya [8], which, in terms of Adams [1], is identical to the set of open subsets with a certain segment property.

Our paper is organized as follows. In Section 2, we give some definitions and preliminary facts for domains of class $\mathcal{C}$. In Section 3, we establish a divergence free partition of vector-valued Sobolev functions mentioned above. In Section 4, we study properties of the function spaces $H_{0,\sigma}^1(\Omega)$ and $H_{0,\sigma}^1(D \setminus \overline{\Omega})$. Finally, in Section 5, we prove the existence theorem of a shape optimization problem for the stationary Navier-Stokes equation.

2. Preliminary facts for open subsets of class $\mathcal{C}$

Let Ω be a bounded open subset in $\mathbf{R}^n$, $n \geq 1$. We say that Ω is of class $\mathcal{C}$ or has property $\mathcal{C}$, if the following properties (2.1), (2.2) and (2.3) are satisfied:

There is a family $\mathcal{F}_\Omega$ of real valued continuous functions g defined over $\overline{S(0, k_\Omega)}$, where $S(0, k_\Omega) \subset \mathbf{R}^{n-1}$ is the open ball centered at 0 and of radius $k_\Omega > 0$, such that

$$\partial\Omega = \bigcup_{g \in \mathcal{F}_\Omega} \{o_g + R_g((s, g(s))); s \in S(0, k_\Omega)\}. \quad (2.1)$$

Here $o_g \in \partial\Omega$ is a certain point which gives the center of the local sys-

tem of axes and $R_g : \mathbf{R}^n \rightarrow \mathbf{R}^n$ is a certain unitary rotation such that $R_g(0, \dots, 0, 1) = y_g$, where the unit vector y_g gives the local vertical direction. Moreover, for any $g \in \mathcal{F}_\Omega$, there is an $a_\Omega > 0$ such that

$$o_g + R_g((s, g(s))) - ty_g \in \Omega, t \in (0, a_\Omega), s \in S(0, k_\Omega), \quad (2.2)$$

$$o_g + R_g((s, g(s))) + ty_g \in \mathbf{R}^n \setminus \overline{\Omega}, t \in (0, a_\Omega), s \in S(0, k_\Omega). \quad (2.3)$$

By the Lebesgue Lemma, it follows from (2.1) that there is an $r_\Omega \in (0, k_\Omega)$ such that the restricted local charts defined on $\overline{S(0, r_\Omega)}$ also give a covering of $\partial\Omega$:

$$\partial\Omega = \bigcup_{g \in \mathcal{F}_\Omega} \{o_g + R_g((s, 0)) + g(s)y_g; s \in \overline{S(0, r_\Omega)}\}. \quad (2.4)$$

The above definition was given in [14], which is a slight modification of the corresponding definition given by Maz'ya [8]. Roughly speaking, Ω is of class $\mathcal{C}$ if $\partial\Omega$ is locally a graph of a continuous function with segment property. We easily see that if Ω is of class $\mathcal{C}$, then there are open sets $\{V_j\}_{j=1}^m$ such that

$$\partial\Omega \subset \bigcup_{j=1}^m V_j, \quad (2.5)$$

$$V_j = \{x \in \mathbf{R}^n; x = p_j(s) + t\mathbf{v}_j, s \in S(0, k_\Omega), t \in (-d, d)\}, \quad (2.6)$$

with $\mathbf{v}_j$ a unit vector, $p_j(s) = o_j + R_j((s, g_j(s)))$, $g_j \in \mathcal{F}_\Omega$, $o_j \in \partial\Omega$ and R_j a unitary rotation satisfying $R_j((0, \dots, 0, 1)) = \mathbf{v}_j$. Moreover, by the segment property in (2.2) and (2.3), we may take $d > 0$ small enough such that for any $j = 1, \dots, m$,

$$V_j^{-1} \equiv \{x \in \mathbf{R}^n; x = p_j(s) + t\mathbf{v}_j, s \in S(0, k_\Omega), t \in (-d, 0)\} \subset \Omega, \quad (2.7)$$

$$V_j^1 \equiv \{x \in \mathbf{R}^n; x = p_j(s) + t\mathbf{v}_j, s \in S(0, k_\Omega), t \in (0, d)\} \subset \mathbf{R}^n \setminus \overline{\Omega}, \quad (2.8)$$

and

$$\{x \in \mathbf{R}^n; x = p_j(s) - d\mathbf{v}_j, s \in S(0, k_\Omega)\} \subset \Omega. \quad (2.9)$$

We also notice that

$$V_j^0 \equiv \{x \in \mathbf{R}^n; x = p_j(s), s \in S(0, k_\Omega)\} \subset \partial\Omega, \quad (2.10)$$

Now let D and B be bounded open subsets in $\mathbf{R}^n$ with $B \subset\subset D$, and let a, r and k be positive constants with $r > k$. Define $\mathcal{O}(a, r, k)$ to be *the family*

of all open subsets contained in B such that every $\Omega \in \mathcal{O}(a, r, k)$ is of class $\mathcal{C}$ with $k_\Omega \geq k > 0, a_\Omega > 0, r_\Omega \leq r$ (where k_Ω, a_Ω and r_Ω are given as in (2.1), (2.2)-(2.3) and (2.4)). In what follows, we will write $\mathcal{O}$ for $\mathcal{O}(a, r, k)$, for simplicity of notation. Moreover, we require that the family of functions $\mathcal{F} = \bigcup_{\Omega \in \mathcal{O}} \mathcal{F}_\Omega$ is equi-continuous and equi-bounded on $S(0, k)$. We define the usual topology on $\mathcal{O}$ by the Hausdorff-Pompeiu distance between the complementary sets (which are closed):

$$\rho(\Omega_1, \Omega_2) = \text{dist}(\overline{B} \setminus \Omega_1, \overline{B} \setminus \Omega_2) \quad \text{for all } \Omega_1, \Omega_2 \in \mathcal{O}. \quad (2.11)$$

Denote by Hlim , the limit in terms of (2.11). The following lemmas will be used to study the existence of our shape optimization problem in section 5.

Lemma 2.1 ([14]) Let $\{\Omega_m\}_{m=1}^\infty \subset \mathcal{O}$. There exists a subsequence $\{\Omega_{m_k}\}_{k=1}^\infty$ of $\{\Omega_m\}_{m=1}^\infty$ such that

$$\text{Hlim}_{k \rightarrow \infty} \Omega_{m_k} = \Omega_0 \in \mathcal{O}.$$

Lemma 2.2 ([10]) (Γ -property for $\mathcal{O}$) Let $\{\Omega_m\}_{m=1}^\infty \subset \mathcal{O}$ and let $\Omega_0 \in \mathcal{O}$ be such that $\text{Hlim}_{m \rightarrow \infty} \Omega_m = \Omega_0$. Then for any open subset K with $K \subset \subset \Omega_0$ (i.e., $\overline{K} \subset \Omega_0$), there is a natural number $m(K)$ such that for all $m \geq m(K)$, $\overline{K} \subset \Omega_m$.

Lemma 2.3 ([15]) ($\hat{\Gamma}$ -property for $\mathcal{O}$) Let $\{\Omega_m\}_{m=1}^\infty \subset \mathcal{O}$ and $\Omega_0 \in \mathcal{O}$ be such that $\text{Hlim}_{m \rightarrow \infty} \Omega_m = \Omega_0$. Then for any open subset K with $\overline{K} \subset \mathbf{R}^n \setminus \Omega_0$, there is a natural number $m(K)$ such that for any $m \geq m(K)$, $\overline{K} \subset D \setminus \Omega_m$.

3. Divergence free partition of vector-valued Sobolev functions —an application of the Hodge theorem

In this section, we shall apply the Hodge theorem to establish a divergence free partition for vector-valued Sobolev functions in $\mathbf{R}^n, n \geq 1$. We will refer the reader to the book of George de Rham [5] for a detailed account on the Hodge Theory.

We recall that for an open subset $\Omega \subset \mathbf{R}^n$, $D_0^1(\Omega) \equiv \{\mathbf{v} \in H_0^1(\Omega)^n; \text{div } \mathbf{v} = 0\}$.

Theorem 3.1 Let D be a bounded open subset in $\mathbf{R}^n$ and let $\mathbf{u} \in D_0^1(D)$. Suppose that $U_i, i = 1, \dots, m$, are bounded open subsets in $\mathbf{R}^n$

such that $\bigcup_{j=1}^m U_j \subset\subset D$. Then for any open subset $\hat{U} \subset\subset \bigcup_{j=1}^m U_j$, there are functions $\{\mathbf{u}_j\}_{j=1}^m$ such that

$$\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j \quad \text{over } \hat{U}$$

and for $j = 1, \dots, m$, $\mathbf{u}_j \in D_0^1(U_j)$ and

$$\|\mathbf{u}_j\|_{H^1(U_j)} \leq C\|\mathbf{u}\|_{H^1(D)},$$

where C is independent of $\mathbf{u} \in D_0^1(D)$.

Proof of Theorem 3.1 Write $\mathbf{u} = (u_1, \dots, u_n)$ and

$$\omega = \sum_{i=1}^n (-1)^{i+1} u_i dx_1 \wedge \dots \wedge \hat{dx}_i \wedge \dots \wedge dx_n. \quad (3.1)$$

Then $d\omega = \operatorname{div} \mathbf{u} \, dx_1 \wedge \dots \wedge dx_n = 0$. Hence, ω is a closed $(n-1)$ -form over $\mathbf{R}^n$ with $H^1(\mathbf{R}^n)$ -regular coefficients and $\operatorname{spt}(\omega) \subset \overline{D}$, where $\operatorname{spt}(\omega)$ denotes the support of ω .

Let S^n be the standard sphere in $\mathbf{R}^{n+1}$ and let g be the standard metric on S^n . We fix a Sobolev space $H^\alpha(S^n)$ of order α for $\alpha \in \mathbf{R}$ by using the metric g and a certain partition of unity over S^n . Let $\pi : S^n \setminus N \rightarrow \mathbf{R}^n$ with $N = (0, \dots, 0, 1)$ be the stereographic projection and let

$$\omega^* = \begin{cases} \pi^*(\omega), & \text{for } p \in S^n \setminus N, \\ 0, & \text{for } p = N. \end{cases}$$

Since ω has compact support in $\mathbf{R}^n$, ω^* is an $(n-1)$ -form over S^n with $H^1(S^n)$ -regular coefficients. Moreover ω^* is identically zero in a small neighborhood of N . Notice that $d\omega^* = 0$.

Let $\Delta_{g,n-1}$ be the standard Betrami-Laplacian operator over S^n acting on the space of $(n-1)$ -forms. Consider the equation

$$\Delta_{g,n-1} \sigma^* = \omega^*. \quad (3.2)$$

By the Hodge theorem, $\operatorname{Ker} \Delta_{g,n-1} \cong H_{deRham}^{n-1}(S^n, \mathbf{R}) = \{0\}$. Hence, equation (3.2) admits a unique solution σ^* with the estimate

$$\|\sigma^*\|_{H^3(S^n)} \leq C\|\omega^*\|_{H^1(S^n)},$$

where C is independent of the choice of ω^* .

Notice that $\Delta_{g,n-1}\sigma^* = dd^*\sigma^* + d^*d\sigma^* = \omega^*$ and $d\omega^* = 0$. Here d^* is the Hilbert space adjoint of the exterior differential operator. It follows that

$$(dd^*\sigma^* + d^*d\sigma^*, d^*d\sigma^*) = (\omega^*, d^*d\sigma^*) = (d\omega^*, d\sigma^*) = 0.$$

Since $(dd^*\sigma^*, d^*d\sigma^*) = (d^2d^*\sigma^*, d\sigma^*) = 0$, we get $d^*d\sigma^* = 0$. Thus $dd^*\sigma^* = \omega^*$. Write $\tau^* = d^*\sigma^*$. Then $d\tau^* = \omega^*$ and

$$\|\tau^*\|_{H^2(S^n)} \leq C\|\omega^*\|_{H^1(S^n)} \leq C\|\omega\|_{H^1(D)}.$$

Here and in what follows, C denotes a constant independent of the choice of ω with $\text{spt}\omega \subset \overline{D}$. However the constant C may be different in different context.

Let $\tau = (\pi^{-1})^*(\tau^*|_{S^n \setminus N})$. Then

$$d\tau = d((\pi^{-1})^*(\tau^*|_{S^n \setminus N})) = (\pi^{-1})^*d(\tau^*|_{S^n \setminus N}) = (\pi^{-1})^*(\omega^*|_{S^n \setminus N}) = \omega,$$

and

$$\|\tau\|_{H^2(D)} \leq C\|\tau^*\|_{H^2(S^n)} \leq C\|\omega\|_{H^1(D)}. \quad (3.3)$$

We next let $\{\chi_j\}_{j=1}^m$ be such that $\chi_j \in C_0^\infty(U_j)$, $0 \leq \chi_j \leq 1$, and $\sum_{j=1}^m \chi_j = 1$ over $\hat{U} \subset \subset \bigcup_{j=1}^m U_j$. Set $\tau_j = \chi_j \tau$. Then $\text{spt}\tau_j \subset U_j$ and

$$\sum_{j=1}^m d\tau_j = \omega \quad \text{over } \hat{U}.$$

Moreover, $d\tau_j$ is an $(n-1)$ -form over $\mathbf{R}^n$ with H^1 -regular coefficients and $\text{spt}(d\tau_j) \subset U_j$. Write for each $j = 1, \dots, m$,

$$d\tau_j = \sum_{i=1}^n (-1)^{i+1} b_{i,j} dx_1 \wedge \dots \wedge \hat{dx}_i \wedge \dots \wedge dx_n, \quad (3.4)$$

where $b_{i,j} \in H_0^1(U_j)$, $i = 1, \dots, n$.

Let $\mathbf{u}_j = (b_{1,j}, \dots, b_{n,j})$. Then $\mathbf{u}_j \in H_0^1(U_j)^n$. Since $0 = d^2\tau_j = (\text{div}\mathbf{u}_j)dx_1 \wedge \dots \wedge dx_n$, it follows that $\text{div}\mathbf{u}_j = 0$. Moreover, by (3.3) we have

$$\|\mathbf{u}_j\|_{H^1(U_j)} \leq C\|\tau_j\|_{H^2(U_j)} \leq C\|\tau\|_{H^2(D)} \leq C\|\omega\|_{H^1(D)} \leq C\|\mathbf{u}\|_{H^1(D)}.$$

On the other hand, since

$$\sum_{j=1}^m d\tau_j = \omega \quad \text{over} \quad \hat{U}$$

it follows from (3.1) and (3.4) that

$$\begin{aligned} & \sum_{j=1}^m \sum_{i=1}^n (-1)^{i+1} b_{i,j} dx_1 \wedge \cdots \wedge \hat{dx}_i \wedge \cdots \wedge dx_n \\ &= \sum_{j=1}^n (-1)^{i+1} u_i dx_1 \wedge \cdots \wedge \hat{dx}_i \wedge \cdots \wedge dx_n \quad \text{over} \quad \hat{U}, \end{aligned}$$

which implies

$$u_i = \sum_{j=1}^m b_{i,j} \quad \text{over} \quad \hat{U}, \quad i = 1, \dots, n.$$

Thus, $\mathbf{u} = \sum_{i=1}^m \mathbf{u}_j$ over $\hat{U}$. This completed the proof. ■

Corollary 3.2 Let $D \subset \mathbf{R}^n$ be a bounded open subset and let $\mathbf{u} \in D_0^1(D)$. Suppose that $\{U_j\}_{j=1}^m$ is an open covering of $\overline{D}$ with $\bigcup_{j=1}^m U_j \subset \subset B_R(0)$, the ball in $\mathbf{R}^n$ centered at 0 and of radius R . Then there are functions $\{\mathbf{u}_j\}_{j=1}^m$, such that

$$\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j \quad \text{over} \quad D$$

and for all $j = 1, \dots, m$, $\mathbf{u}_j \in D_0^1(U_j)$ and $\|\mathbf{u}_j\|_{H^1(U_j)} \leq C\|\mathbf{u}\|_{H^1(D)}$. Here C is independent of $\mathbf{u} \in D_0^1(D)$, but depends on R .

Proof of Corollary 3.2 Let $\omega, \pi, S^n, N, \omega^*, \sigma^*, \tau^*$ and τ be given as in the proof of theorem 3.1. We then have $d\tau = \omega$ and $\|\tau^*\|_{H^2(S^n)} \leq C\|\omega\|_{H^1(D)}$. Let $\chi_{B_R} \in C_0^\infty(B_R)$ be a cutting function with $\chi_{B_R} = 1$ over $\bigcup_{j=1}^m U_j$. Then

$$d(\chi_{B_R} \tau) = \omega \quad \text{over} \quad \bigcup_{j=1}^m U_j$$

and

$$\begin{aligned} \|\chi_{B_R} \tau\|_{H^2(\mathbf{R}^n)} &\leq C\|\pi^*(\chi_{B_R} \tau)\|_{H^2(S^n)} \leq C\|\tau^*\|_{H^2(S^n)} \\ &\leq C\|\omega^*\|_{H^1(S^n)} \leq C\|\omega\|_{H^1(D)}. \end{aligned} \quad (3.5)$$

Now let $\{\chi_j\}_{j=1}^m$ be such that $0 \leq \chi_j \leq 1, \chi_j \in C_0^\infty(U_j), \sum_{j=1}^m \chi_j = 1$ over D . Then

$$d(\chi_{B_R} \tau \sum_{j=1}^m \chi_j) = \omega \quad \text{over } D.$$

Let $\tau_j = \chi_{B_R} \chi_j \tau$. Then $\text{spt} \tau_j \subset U_j$. Thus it follows from (3.5) that $d\tau_j$ is an $(n-1)$ -form with $H^1(\mathbf{R}^n)$ -smooth coefficients and

$$\|d\tau_j\|_{H^1(\mathbf{R}^n)} \leq C\|\omega\|_{H^1(D)}.$$

Write

$$d\tau_j = \sum_{i=1}^n (-1)^{i+1} b_{i,j} dx_1 \wedge \cdots \wedge \hat{dx}_i \wedge \cdots \wedge dx_n, j = 1, \dots, m,$$

and let $\mathbf{u}_j = (b_{1,j}, \dots, b_{n,j})$ for $j = 1, \dots, m$. Then by the same argument as that in the proof of Theorem 3.1, we get

$$\begin{aligned} \mathbf{u} &= \sum_{j=1}^m \mathbf{u}_j \quad \text{over } D, \\ \mathbf{u}_j &\in D_0^1(U_j), j = 1, \dots, m, \end{aligned}$$

and

$$\|\mathbf{u}_j\|_{H^1(U_j)} \leq C\|\mathbf{u}\|_{H^1(D)}, j = 1, \dots, m.$$

This completes the proof. ■

4. A localization property of the divergence free decomposition

In this section, we shall prove a localization property of the divergence free decomposition established in section 3. Based on it, we obtain some identities for certain fundamental function spaces related to the Navier-Stokes equations.

Theorem 4.1 Let D be a bounded open subset in $\mathbf{R}^2$, and let $\Omega = \bigcup_{j=1}^N \Omega_j \subset\subset D$ be a finite union of open and connected subsets in $\mathbf{R}^2$ with $\overline{\Omega_i} \cap \overline{\Omega_j} = \emptyset$ for $i \neq j$. Assume that $\{U_j\}_{j=1}^m$ be an open covering of $\partial\Omega$ in $\mathbf{R}^2$ with $\bigcup_{j=1}^m U_j \subset\subset D$. Let $\mathbf{u} \in D_0^1(D)$ with $\mathbf{u} = 0$ over Ω . Then there are functions $\{\mathbf{u}_j\}_{j=1}^m$ such that

$$\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j$$

over a certain neighborhood $\hat{U}$ of $\partial\Omega$ in $\mathbf{R}^2$ with $\hat{U} \subset \subset \bigcup_{j=1}^m U_j$, and for all $j = 1, \dots, m$, $\mathbf{u}_j \in D_0^1(U_j)$, $\mathbf{u}_j = 0$ over Ω and

$$\|\mathbf{u}_j\|_{H^1(U_j)} \leq C\|\mathbf{u}\|_{H^1(D)},$$

where C is a positive constant independent of $\mathbf{u} \in D_0^1(D)$.

Proof of Theorem 4.1 Step 1. We shall first prove the theorem in the case that Ω is connected.

Let ω, ω^*, τ^* and π be given as in the proof of Theorem 3.1. We then have $\omega^* = d\tau^*$ and

$$\|\tau^*\|_{H^2(S^2)} \leq C\|\omega\|_{H^1(S^2)} \leq C\|\omega\|_{H^1(D)}. \quad (4.1)$$

Since $\mathbf{u} = 0$ over Ω , it follows that $\omega = 0$ over Ω , which implies that $\omega^* = 0$ over $\pi^{-1}(\Omega)$. Hence, $d\tau^* = 0$ over $\pi^{-1}(\Omega)$. Because τ^* is a 0-form and Ω is connected, there is a constant $c_0 \in \mathbf{R}$ such that

$$\tau^* = c_0 \quad \text{over} \quad \pi^{-1}(\Omega). \quad (4.2)$$

By (4.1) and (4.2), we obtain

$$|c_0| \leq C\|\omega\|_{H^1(D)}. \quad (4.3)$$

Let $\tilde{\tau}^* = \tau^* - c_0$. Then $\tilde{\tau}^* = 0$ over $\pi^{-1}(\Omega)$ and $d\tilde{\tau}^* = d(\tau^* - c_0) = d\tau^* = \omega^*$, moreover, it follows from (4.1) and (4.3) that

$$\|\tilde{\tau}^*\|_{H^2(S^2)} \leq C\|\omega\|_{H^1(D)}.$$

Let $\tilde{\tau} = (\pi^{-1})^*(\tilde{\tau}^*|_{S^2 \setminus N})$. Then $\tilde{\tau} = 0$ over Ω , $d\tilde{\tau} = \omega$ and

$$\|\tilde{\tau}\|_{H^2(D)} \leq C\|\pi^*\tilde{\tau}\|_{H^2(S^2)} \leq C\|\tilde{\tau}^*\|_{H^2(S^2)} \leq C\|\omega\|_{H^1(D)}. \quad (4.4)$$

Let $\hat{U}$ be any neighborhood of $\partial\Omega$ such that $\hat{U} \subset \subset \bigcup_{j=1}^m U_j$ and let $\{\chi_j\}_{j=1}^m$ be such that $0 \leq \chi_j \leq 1$, $\chi_j \in C_0^1(U_j)$ and $\sum_{j=1}^m \chi_j = 1$ over $\hat{U}$. Then we have

$$d\left(\sum_{j=1}^m \chi_j \tilde{\tau}\right) = \omega \quad \text{over} \quad \hat{U}.$$

Let $\tilde{\tau}_j = \chi_j \tilde{\tau}$, $j = 1, \dots, m$. Then $\tilde{\tau}_j \in H_0^1(U_j)$, $\tilde{\tau}_j = 0$ over Ω and

$$\sum_{j=1}^m d\tilde{\tau}_j = \omega \quad \text{over} \quad \hat{U}.$$

Write, for $j = 1, \dots, m$, $d\tilde{\tau}_j = b_{1,j}dx_2 - b_{2,j}dx_1$ for certain $b_{1,j}, b_{2,j} \in H_0^1(U_j)$ with $b_{i,j} = 0$ over Ω . Let $\mathbf{u}_j = (b_{1,j}, b_{2,j})$, $j = 1, \dots, m$. Then one can easily check that for all $j = 1, \dots, m$, $\mathbf{u}_j = 0$ over Ω , $\mathbf{u}_j \in D_0^1(U_j)$ and $\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j$ over $\hat{U}$. Moreover, it follows from (4.4) that

$$\begin{aligned} \|\mathbf{u}_j\|_{H^1(U_j)} &\leq C(\|b_{1,j}\|_{H^1(U_j)} + \|b_{2,j}\|_{H^1(U_j)}) \\ &\leq C\|d\tilde{\tau}_j\|_{H^1(U_j)} \leq C\|\tilde{\tau}\|_{H^2(D)} \\ &\leq C\|\omega\|_{H^1(D)} \leq C\|\mathbf{u}\|_{H^1(D)}. \end{aligned}$$

This proves the theorem in case Ω is connected.

Step 2. We next prove the theorem in case Ω is multiple-connected. For brevity of notation, we assume that Ω is 2-connected. (The general case can be similarly done through an induction argument.) Then $\Omega = \Omega_1 \cup \Omega_2$ with $\overline{\Omega_1} \cap \overline{\Omega_2} = \emptyset$, Ω_1 and Ω_2 being connected.

By refining the covering $\{U_j\}_{j=1}^m$ of $\partial\Omega$, we may assume, without loss of generality, that there are open subsets $\{U_{j,1}\}_{j=1}^m$ and $\{U_{j,2}\}_{j=1}^m$ such that

$$\begin{aligned} \bigcup_{j=1}^m U_{j,1} &\supset \partial\Omega_1, \quad \bigcup_{j=1}^m U_{j,2} \supset \partial\Omega_2, \\ U_{j,i} &\subset U_j \quad \text{for } j = 1, \dots, m, \quad i = 1, 2, \\ U_{l,1} \cap U_{j,2} &= \emptyset \quad \text{for } l, j = 1, \dots, m. \end{aligned}$$

Let $\hat{U}_1$ and $\hat{U}_2$ be neighborhoods of $\partial\Omega_1$ and $\partial\Omega_2$, respectively, such that

$$\hat{U}_1 \subset \subset \bigcup_{j=1}^{m_1} U_{j,1}, \quad \hat{U}_2 \subset \subset \bigcup_{j=1}^{m_2} U_{j,2}.$$

Since $\mathbf{u} \in D_0^1(D)$ and $\mathbf{u} = 0$ over Ω_1 , it follows from Step 1 that there are functions $\{\mathbf{u}_{j,1}\}_{j=1}^m$ such that $\mathbf{u} = \sum_{j=1}^m \mathbf{u}_{j,1}$ over $\hat{U}_1$ and for each $j = 1, \dots, m$, $\mathbf{u}_{j,1} \in D_0^1(U_{j,1})$, $\mathbf{u}_{j,1} = 0$ over Ω_1 , and

$$\|\mathbf{u}_{j,1}\|_{H^1(U_{j,1})} \leq C\|\mathbf{u}\|_{H^1(D)}.$$

Notice that for $j = 1, \dots, m$, $\mathbf{u}_{j,1} = 0$ over Ω_2 , for $\Omega_2 \cap U_{j,1} = \emptyset$ and $\mathbf{u}_{j,1} \in D_0^1(U_{j,1})$.

Now let $\mathbf{v} = \mathbf{u} - \sum_{j=1}^m \mathbf{u}_{j,1}$. Then $\mathbf{v} \in D_0^1(D)$ and $\mathbf{v} = 0$ over Ω_2 . Again, by Step 1, there are functions $\{\mathbf{u}_{j,2}\}_{j=1}^m$ such that for each $j = 1, \dots, m$, $\mathbf{u}_{j,2} \in D_0^1(U_{j,2})$, $\mathbf{u}_{j,2} = 0$ over Ω_2 ,

$$\|\mathbf{u}_{j,2}\|_{H^1(U_{j,2})} \leq C\|\mathbf{v}\|_{H^1(D)} \leq C\|\mathbf{u}\|_{H^1(D)},$$

and $\mathbf{v} = \sum_{l=1}^m \mathbf{u}_{l,2}$ over $\hat{U}_2$. Notice also that for $j = 1, \dots, m$, $\mathbf{u}_{j,2} = 0$ over Ω_1 , for $U_{j,2} \cap \Omega_1 = \emptyset$ and $\mathbf{u}_{j,2} \in D_0^1(U_{j,2})$.

Finally, let $\mathbf{u}_l = \mathbf{u}_{l,1} + \mathbf{u}_{l,2}$, $l = 1, \dots, m$. Then it is clear that

$$\mathbf{u} = \sum_{l=1}^m \mathbf{u}_l \quad \text{over} \quad \hat{U} \equiv \hat{U}_1 \bigcup \hat{U}_2,$$

and for all $l = 1, \dots, m$, $\mathbf{u}_l \in D_0^1(U_l)$, $\mathbf{u}_l = 0$ over Ω and $\|\mathbf{u}_l\|_{H^1(U_l)} \leq C\|\mathbf{u}\|_{H^1(D)}$. This completes the proof. ■

By the almost same argument as above with some slight modification, we can get the following.

Theorem 4.1' Let D be a bounded open subset in $\mathbf{R}^2$, and let $\Omega = \bigcup_{j=1}^N \Omega_j \subset\subset D$ be a finite union of open and connected subsets with $\overline{\Omega_i} \cap \overline{\Omega_j} = \emptyset$ for $i \neq j$. Suppose that Γ is a compact subset of $\partial\Omega$ and $\{U_j\}_{j=1}^m$ is an open covering of Γ in $\mathbf{R}^2$ with $\bigcup_{j=1}^m U_j \subset\subset D$. Let $\mathbf{u} \in D_0^1(D)$ with $\mathbf{u} = 0$ over Ω . Then there are functions $\{\mathbf{u}_j\}_{j=1}^m$ such that

$$\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j$$

over a certain neighborhood $\hat{U}$ of Γ in $\mathbf{R}^2$ with $\hat{U} \subset\subset \bigcup_{j=1}^m U_j$, and for all $j = 1, \dots, m$, $\mathbf{u}_j \in D_0^1(U_j)$, $\mathbf{u}_j = 0$ over Ω and

$$\|\mathbf{u}_j\|_{H^1(U_j)} \leq C\|\mathbf{u}\|_{H^1(D)},$$

where C is a positive constant independent of $\mathbf{u} \in D_0^1(D)$.

We next prove a version of Theorem 4.1 for domains in $\mathbf{R}^3$. To this aim, we need to recall some results from Algebraic Topology. Two good references for the topological results which we will quote here are [9] and [2].

Suppose that Ω is a bounded open subset in $\mathbf{R}^n$, $n \geq 1$, with $\partial\Omega$ a compact topological manifold of dimension $n-1$. By making use of results in Algebraic Topology (see pp.227, (26.17.8), [9]), there is a bounded domain $\Omega' \subset\subset \mathbf{R}^n$ with smooth boundary $\partial\Omega'$ such that $\Omega' \supset\supset \Omega$ and Ω' can be retracted to Ω . Namely, there is a continuous map $\gamma : \Omega' \rightarrow \overline{\Omega}$ with $\gamma|_{\overline{\Omega}} = id$ (the identity map). This implies that the homological group $H_1(\Omega, \mathbf{R})$ is of finite dimension and can be naturally embedded as a vector space of $H_1(\Omega', \mathbf{R})$. Let $\{[\gamma_1], \dots, [\gamma_k]\}$ be a basis of $H_1(\Omega, \mathbf{R})$ with $\gamma_j :$

$[0, 1] \rightarrow \Omega$ a smooth closed Jordan curve for each j (see pp.63, (12.1), [9]). We can then extend $\{[\gamma_1], \dots, [\gamma_k]\}$ to a basis $\{[\gamma_1], \dots, [\gamma_k], \dots, [\gamma_M]\}$ of $H_1(\Omega', \mathbf{R})$. Let $\{[\tau_1], \dots, [\tau_M]\}$ be the dual basis of $\{[\gamma_1], \dots, [\gamma_k], \dots, [\gamma_M]\}$ in $H_{deRham}^1(\Omega', \mathbf{R})$. Then $\{[\tau_1], \dots, [\tau_k]\}$, when restricted to Ω , gives a dual basis of $\{[\gamma_1], \dots, [\gamma_k]\}$, namely, $\int_{\gamma_i} \tau_j = \delta_j^i$. Here $\{\tau_j\}$ are smooth 1-forms over Ω' .

Lemma 4.2 Under the above notations, further assume that $\Omega \subset \mathbf{R}^3$ is connected and of class $\mathcal{C}$. Let τ be a 1-form over Ω' with $H^2(\Omega')$ -regular coefficients. Also assume that $d\tau = 0$ over Ω . Then there exists $h \in H_0^2(\Omega')$ and $\{c_j\}_{j=1}^k \subset \mathbf{R}$ such that

$$\tau = dh + \sum_{j=1}^k c_j \tau_j \quad \text{over } \Omega,$$

and

$$\|h\|_{H^2(\Omega')} \leq C \|\tau\|_{H^2(\Omega')}, \quad |c_j| \leq C \|\tau\|_{H^2(\Omega')}, \quad j = 1, \dots, k.$$

Here C is a positive constant depending only on Ω and Ω' .

Proof of Lemma 4.2 Let $\psi_0 \in C_0^\infty(-1, 1)$ be such that $\psi_0 \geq 0$ and $\int_{\mathbf{R}^3} \psi(x) dx = 1$ where $\psi(x) \equiv \psi_0(|x|^2)$, and let $\tau_\delta(x) = \int_{\mathbf{R}^3} \tau(x-y) \psi_\delta(y) dy$, where $\delta > 0$ and $\psi_\delta(y) = \delta^{-3} \psi(\delta^{-1} y)$. Then by the Friedrich smoothing lemma, for any $\delta_0 \ll 1$, $\tau_\delta \rightarrow \tau$ in $H^2(\Omega'_{\delta_0})$ as $\delta \rightarrow 0$ where $\Omega'_{\delta_0} \equiv \{x \in \Omega'; \text{dist}(x, \partial\Omega') > \delta_0\}$. Moreover, $\tau_\delta \in \Lambda^1(\Omega'_\delta)$, i.e., τ_δ is a smooth 1-form over Ω'_δ and $d\tau_\delta = 0$ over Ω_δ , where $\Omega_\delta \equiv \{x \in \Omega; \text{dist}(x, \partial\Omega) > \delta\}$.

Next, since τ has $H^2(\Omega')$ -regular coefficients, for each piecewise smooth curve γ in Ω' , the restriction of τ on γ is well-defined and is L^2 -integrable by the trace theorem. Hence, we can well define $\int_\gamma \tau$, moreover, by the trace theorem,

$$|\int_\gamma \tau| \leq C(\gamma) \|\tau\|_{H^1(\Omega')},$$

where $C(\gamma)$ depends on γ .

Now let $c_j = \int_{\gamma_j} \tau$, $j = 1, \dots, k$, and $\hat{\tau} = \tau - \sum_{j=1}^k c_j \tau_j$. It is clear that

$$|c_j| \leq C(\gamma_j) \|\tau\|_{H^1(\Omega')}.$$

We claim that for each piecewise smooth curve $\gamma : [0, 1] \rightarrow \Omega$, $\int_\gamma \hat{\tau}$ depends only on $\gamma(0)$ and $\gamma(1)$.

Indeed, for each $\gamma' : [0, 1] \rightarrow \Omega$ piecewise smooth curve with $\gamma'(0) = \gamma(0), \gamma'(1) = \gamma(1)$, since $\gamma' - \gamma$ is homotopical to $\sum_{j=1}^k d_j \gamma_j$ for a certain choice of $\{d_j\}_{j=1}^k \subset \mathbf{R}$, we get

$$\begin{aligned} \int_{\gamma' - \gamma} (\tau_\delta - \sum_{i=1}^k c_i \tau_i) &= \sum_{j=1}^k \int_{\gamma_j} d_j (\tau_\delta - \sum_{i=1}^k c_i \tau_i) \\ &= \sum_{i,j=1}^k (d_j \int_{\gamma_j} \tau_\delta - c_i d_j \delta_i^j) \\ &= \sum_{j=1}^k d_j (\int_{\gamma_j} \tau_\delta - c_j). \end{aligned}$$

Letting $\delta \rightarrow 0$, we get

$$\int_{\gamma' - \gamma} (\tau - \sum_{i=1}^k c_i \tau_i) = \sum_{j=1}^k d_j (c_j - c_j) = 0.$$

Now we fix $p_0 \in \Omega$. Since Ω is connected, for any $p \in \Omega$, there is a piecewise smooth curve $\gamma : [0, 1] \rightarrow \Omega$ with $\gamma(0) = p_0, \gamma(1) = p$. We define $h : \Omega \rightarrow \mathbf{R}$ by

$$h(p) = \int_{\gamma} (\tau - \sum_{j=1}^k c_j \tau_j) = \int_{\gamma} \hat{\tau}.$$

It is clear that h is well-defined over Ω . By the standard arguments in calculus, one can easily verify that $dh = \hat{\tau}$ over Ω and thus $h \in H^3(\Omega)$. Moreover,

$$\|h\|_{H^3(\Omega)} \leq C \|\tau\|_{H^2(\Omega')}.$$

Through the proof, as mentioned before, C denotes a positive constant depending only on Ω and Ω' , which may be different in different context.

Next, we are going to extend h from Ω to Ω' so that the extension is the desired function. To this end, we observe first that since Ω is of class $\mathcal{C}$, there exist open sets $\{V_j\}_{j=1}^m$ such that (2.5)-(2.10) are fulfilled. Moreover, by taking d small enough, we may have

$$\bigcup_{i=1}^m V_i \subset \subset \Omega'.$$

Now we define the extension of h on V_j as follows:

$$\tilde{h}_j(p_j(s) + t\mathbf{v}_j) = \int_{C_{p_j(s)+t\mathbf{v}_j}} \hat{\tau} + h(p_j(s) - d\mathbf{v}_j),$$

for $s \in S(0, k_\Omega)$ and $t \in (-d, d)$, where $C_{p_j(s)+t\mathbf{v}_j}$ denotes the segment connecting $p_j(s) - d\mathbf{v}_j$ to $p_j(s) + t\mathbf{v}_j$. It is clear that $\tilde{h}_j$ is well-defined. By the trace theorem and a direct computation, we have $\tilde{h}_j \in H^2(U_j)$ with $\|\tilde{h}_j\|_{H^2(U_j)} \leq C\|\tau\|_{H^2(\Omega')}$. Moreover, by (2.5)-(2.10), the whole segment $C_{p_j(s)+t\mathbf{v}_j}$ lies in Ω if $p_j(s) + t\mathbf{v}_j \in \Omega \cap V_j$. Thus for $p_j(s) + t\mathbf{v}_j \in \Omega \cap V_j$,

$$\tilde{h}_j(p_j(s) + t\mathbf{v}_j) = \int_{C_{p_j(s)+t\mathbf{v}_j}} dh + h(p_j(s) - d\mathbf{v}_j) = h(p_j(s) + t\mathbf{v}_j),$$

i.e., $\tilde{h}_j = h$ over $\Omega \cap V_j$.

Let $\hat{U}$ be a neighborhood of $\partial\Omega$ such that $\hat{U} \subset\subset \bigcup_{j=1}^m V_j$ and $\{\chi_j\}_{j=1}^m$ be such that $\chi_j \in C_0^\infty(V_j)$, $0 \leq \chi_j \leq 1$ for $j = 1, \dots, m$, and $\sum_{j=1}^m \chi_j = 1$ over $\hat{U}$. Set

$$\tilde{\tilde{h}}_j(x) = \begin{cases} \tilde{h}_j(x), & x \in V_j, \\ 0, & x \in \Omega' \setminus \overline{V_j}. \end{cases}$$

Then $\chi_j \tilde{\tilde{h}}_j \in H^2(\Omega')$ and

$$\|\chi_j \tilde{\tilde{h}}_j\|_{H^2(\Omega')} \leq C\|\tau\|_{H^2(\Omega')}$$

Let $\tilde{h} = \sum_{j=1}^m \chi_j \tilde{\tilde{h}}_j$, then $\tilde{h}(x) = h(x)$ over $\Omega \cap \hat{U}$ and

$$\|\tilde{h}\|_{H^2(\Omega')} \leq C\|\tau\|_{H^2(\Omega')}$$

Finally, we set

$$\hat{h}(x) = \begin{cases} h(x), & x \in \Omega, \\ \tilde{h}(x), & x \in \hat{U} \cup (\Omega' \setminus \overline{\Omega}). \end{cases}$$

Then one can easily verify that $\hat{h} \in H^2(\Omega')$, $\hat{h} = h$ over Ω and

$$\|\hat{h}\|_{H^2(\Omega')} \leq C\|\tau\|_{H^2(\Omega')}.$$

This completes the proof of Lemma 4.2. ■

Remark 4.3 Let $D \subset \mathbf{R}^n$ be an open subset such that $\Omega \subset\subset D$. Then Ω' in the above lemma can be taken as $\Omega' \subset\subset D$.

Theorem 4.4 Let D be a bounded open subset in $\mathbf{R}^3$ and let $\Omega \subset\subset D$ be of class $\mathcal{C}$. Assume that $\Omega' \supset\supset \Omega$ is given as in Lemma 4.2 such that

$\Omega' \subset\subset D$. Let $\{U_j\}_{j=1}^m$ be an open covering of $\partial\Omega$ in $\mathbf{R}^3$ with $\bigcup_{j=1}^m U_j \subset\subset \Omega'$ and let $\mathbf{u} \in D_0^1(D)$ with $\mathbf{u} = 0$ over Ω . Then there are functions $\{\mathbf{u}_j\}_{j=1}^m$ such that

$$\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j$$

over a certain neighborhood $\hat{U}$ of $\partial\Omega$ in $\mathbf{R}^3$ with $\hat{U} \subset\subset \bigcup_{j=1}^m U_j$, and for all $j = 1, \dots, m$, $\mathbf{u}_j \in D_0^1(U_j)$, $\mathbf{u}_j = 0$ over Ω and

$$\|\mathbf{u}_j\|_{H^1(U_j)} \leq C\|\mathbf{u}\|_{H^1(D)},$$

where C is a positive constant independent of $\mathbf{u} \in D_0^1(D)$.

Proof of Theorem 4.4 Since Ω is of class $\mathcal{C}$, we have $\Omega = \bigcup_{j=1}^N \Omega_j$ with Ω_i being open and connected and $\overline{\Omega_i} \cap \overline{\Omega_j} = \emptyset$ for $i \neq j$. We shall prove the theorem only in the case when Ω is connected. The rest follows from the identical argument as that in Step 2 of the proof of Theorem 4.1.

Let $\omega, \pi, \omega^*, \sigma^*, \tau^*$ and τ be given as in the proof of Theorem 3.1. We have

$$d\tau = \omega \quad \text{and} \quad \|\tau\|_{H^2(D)} \leq C\|\omega\|_{H^1(D)}.$$

Here and throughout the proof, C denotes a positive constant independent of ω , whenever $spt\omega \subset \overline{D}$. However C may be different in different context. Thus $\tau|_{\Omega'}$, the restriction of τ over Ω' , is an 1-form over Ω' with $H^2(\Omega')$ -regular coefficients.

Since $\mathbf{u} = 0$ over Ω , it follows that $d\tau = \omega = 0$ over Ω . Then by Lemma 4.2, there are $h \in H_0^2(\Omega')$, $\{\tau_j\}_{j=1}^k \subset \Lambda^1(\Omega')$ and $\{c_j\}_{j=1}^k \subset \mathbf{R}$ such that

$$\tau = dh + \sum_{j=1}^k c_j \tau_j \quad \text{over } \Omega,$$

$$\|h\|_{H^2(\Omega')} + \sum_{j=1}^k |c_j| \leq C\|\tau\|_{H^2(\Omega')} \leq C\|\tau\|_{H^2(D)}.$$

Let $\hat{U}$ be a neighborhood of $\partial\Omega$ in $\mathbf{R}^3$ such that $\hat{U} \subset\subset \bigcup_{j=1}^m U_j$ and $\{\chi_j\}_{j=1}^m$ be such that $\chi_j \in C_0^\infty(U_j)$, $0 \leq \chi_j \leq 1$ and $\sum_{j=1}^m \chi_j = 1$ over $\hat{U}$.

Let

$$\tau^j = \chi_j(\tau - dh - \sum_{i=1}^k c_i \tau_i), \quad j = 1, \dots, m.$$

Then τ^j is a 1-form over $\mathbf{R}^3$ and

$$d\tau^j = d\chi_j \wedge (\tau - dh - \sum_{i=1}^k c_i \tau_i) + \chi_j d\tau.$$

It is obvious that

$$\sum_{j=1}^m d\tau^j = \omega \quad \text{over} \quad \hat{U},$$

and for all $j = 1, \dots, m$, $d\tau^j \in H_0^1(U_j)$, $d\tau^j = 0$ over $U_j \cap \Omega$. Moreover, it follows from the estimate in Lemma 4.2 that for all $j = 1, \dots, m$,

$$\|d\tau^j\|_{H^1(U_j)} \leq C\|\tau\|_{H^2(D)} \leq C\|\omega\|_{H^1(D)}.$$

Now we write

$$d\tau^j = \sum_{i=1}^3 (-1)^{i+1} b_{i,j} dx_1 \wedge \dots \wedge \hat{dx}_i \wedge \dots \wedge dx_3, \quad j = 1, \dots, m,$$

and let $\mathbf{u}_j = (b_{1,j}, b_{2,j}, b_{3,j})$, $j = 1, \dots, m$. Then one can check easily that

$$\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j \quad \text{over} \quad \hat{U},$$

and for all $j = 1, \dots, m$, $\mathbf{u}_j \in D_0^1(U_j)$, $\mathbf{u}_j = 0$ over Ω and

$$\|\mathbf{u}_j\|_{H^1(U_j)} \leq C\|\mathbf{u}\|_{H^1(D)}.$$

This completes the proof of Theorem 4.4. ■

By the same argument as above with some slight modification, we can get the following.

Theorem 4.4' Let D be a bounded open subset in $\mathbf{R}^3$ and let $\Omega \subset\subset D$ be of class $\mathcal{C}$. Assume that $\Omega' \supset\supset \Omega$ is given as in Lemma 4.2 such that $\Omega' \subset\subset D$. Let Γ be a compact subset of $\partial\Omega$ and let $\{U_j\}_{j=1}^m$ be an open covering of Γ in $\mathbf{R}^3$ with $\bigcup_{j=1}^m U_j \subset\subset \Omega'$. Suppose that $\mathbf{u} \in D_0^1(D)$ with $\mathbf{u} = 0$ over Ω . Then there are functions $\{\mathbf{u}_j\}_{j=1}^m$ such that

$$\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j$$

over a certain neighborhood $\hat{U}$ of Γ in $\mathbf{R}^3$ with $\hat{U} \subset \subset \bigcup_{j=1}^m U_j$, and for all $j = 1, \dots, m$, $\mathbf{u}_j \in D_0^1(U_j)$, $\mathbf{u}_j = 0$ over Ω and

$$\|\mathbf{u}_j\|_{H^1(U_j)} \leq C\|\mathbf{u}\|_{H^1(D)},$$

where C is a positive constant independent of $\mathbf{u} \in D_0^1(D)$.

We next apply the above results to obtain some characterizations for some Sobolev function spaces.

Theorem 4.5 Let $D \subset \mathbf{R}^n$, $n = 2, 3$, be a bounded open Lipschitz subset and let Ω be of class $\mathcal{C}$ such that $\Omega \subset \subset D$. Then $H_{0,\sigma}^1(D \setminus \overline{\Omega}) = \{\mathbf{v} \in H_{0,\sigma}^1(D); \mathbf{v} = 0 \text{ a.e. in } \Omega\}$

Proof of Theorem 4.5 It is clear that $H_{0,\sigma}^1(D \setminus \overline{\Omega}) \subset \{\mathbf{v} \in H_{0,\sigma}^1(D); \mathbf{v} = 0 \text{ a.e. in } \Omega\}$.

Let $\mathbf{u} \in H_{0,\sigma}^1(D)$ satisfying $\mathbf{u} = 0$ a.e. in Ω . Let Ω' be given as in Lemma 4.2 such that $\Omega \subset \subset \Omega' \subset \subset D$. Let $\{V_j\}_{j=1}^m$ be the covering of $\partial\Omega$ as given by (2.5) and (2.6). By taking d in (2.6) sufficiently small, we have

$$\bigcup_{j=1}^m V_j \subset \subset \Omega'.$$

Then by Theorem 4.1 and Theorem 4.4, there are functions $\{\mathbf{u}_j\}_{j=1}^m$ and a neighborhood $\hat{U}$ of $\partial\Omega$ in $\mathbf{R}^n$ such that

$$\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j \quad \text{over } \hat{U}$$

and for all $j = 1, \dots, m$, $\mathbf{u}_j \in D_0^1(V_j)$, $\mathbf{u}_j = 0$ over Ω .

Since $V_j \subset \subset D$ and $\mathbf{u}_j \in D_0^1(V_j)$, it holds that $\mathbf{u}_j \in D_0^1(D)$. Since D is Lipschitz, we have $D_0^1(D) = H_{0,\sigma}^1(D)$ (see Lemma 1.2.1, Chapter 3, [11]). Hence, $\mathbf{u}_j \in H_{0,\sigma}^1(D)$ for $j = 1, \dots, m$.

Now let $\mathbf{u}_0 = \mathbf{u} - \sum_{j=1}^m \mathbf{u}_j$. Then $\mathbf{u}_0 \in H_{0,\sigma}^1(D)$, $\mathbf{u}_0 = 0$ a.e. in $\Omega \cup \hat{U}$. Since $\hat{U} \cup \Omega \supset \supset \Omega$, there exists an open Lipschitz subset $\hat{V}$ such that $\hat{U} \cup \Omega \supset \supset \hat{V} \supset \supset \Omega$. Hence, $\mathbf{u}_0 = 0$ a.e. in $\hat{V}$ which implies by the trace theorem that $\mathbf{u}_0 \in H_0^1(D \setminus \widehat{\hat{V}})^n$. Since $D \setminus \widehat{\hat{V}}$ is Lipschitz and $\text{div} \mathbf{u}_0 = 0$, we have $\mathbf{u}_0 \in H_{0,\sigma}^1(D \setminus \widehat{\hat{V}})$ (see Lemma 1.2.1, Chapter 3, [11]). Thus $\mathbf{u}_0 \in H_{0,\sigma}^1(D \setminus \overline{\Omega})$.

We next claim that $\mathbf{u}_j \in H_{0,\sigma}^1(D \setminus \overline{\Omega})$ for each $j = 1, \dots, m$. To this aim, we fix $j \in \{1, \dots, m\}$ and define $\mathbf{u}_j^t$ for $t \in (0, r)$, where $r = \frac{1}{4} \min(d, \text{dist}(spt \mathbf{u}_j, \partial V_j))$, by $\mathbf{u}_j^t(x) = \mathbf{u}_j(x - t\mathbf{v}_j)$. (We recall that $\mathbf{v}_j$ is given by (2.6).) One can easily check that $\text{div} \mathbf{u}_j^t = 0$ for each $t \in (0, r)$. Now we prove that $\mathbf{u}_j^t \in H_{0,\sigma}^1(D \setminus \overline{\Omega})$ for each $t \in (0, r)$. Indeed, for each fixed $t \in (0, r)$, $spt \mathbf{u}_j^t = spt \mathbf{u}_j + t\mathbf{v}_j$ and $\mathbf{u}_j^t = 0$ over $\Omega + t\mathbf{v}_j$. Thus $spt \mathbf{u}_j^t \subset \subset V_j \setminus \overline{\Omega}$. Therefore, there exists an open Lipschitz subset $U_{j,t} \subset \mathbf{R}^n$ such that $spt \mathbf{u}_j^t \subset \subset U_{j,t} \subset \subset V_j \setminus \overline{\Omega}$. It is clear that $\mathbf{u}_j^t \in H_0^1(U_{j,t})^n$ since $U_{j,t}$ is Lipschitz. This implies that $\mathbf{u}_j^t \in D_0^1(U_{j,t})$, for $\text{div} \mathbf{u}_j^t = 0$. Thus $\mathbf{u}_j^t \in H_{0,\sigma}^1(U_{j,t})$ (see Lemma 1.2.1, Chapter 3, [11]). Hence, $\mathbf{u}_j^t \in H_{0,\sigma}^1(D \setminus \overline{\Omega})$. On the other hand, one can easily verify that

$$\|\mathbf{u}_j^t - \mathbf{u}_j\|_{H^1(D \setminus \overline{\Omega})} \rightarrow 0 \quad \text{as } t \rightarrow 0^+.$$

Thus we have proved that $\mathbf{u}_j \in H_{0,\sigma}^1(D \setminus \overline{\Omega})$.

Since $\mathbf{u} = \sum_{j=0}^m \mathbf{u}_j$, it follows that $\mathbf{u} \in H_{0,\sigma}^1(D \setminus \overline{\Omega})$. This thus completes the proof of Theorem 4.5. ■

Theorem 4.6 Let $\Omega \subset \mathbf{R}^n$ for $n = 2, 3$ be of class $\mathcal{C}$. Then

$$D_0^1(\Omega) = H_{0,\sigma}^1(\Omega).$$

Proof of Theorem 4.6 It suffices to show that $D_0^1(\Omega) \subset H_{0,\sigma}^1(\Omega)$. Let $\mathbf{u} \in D_0^1(\Omega)$ and let D, D_1 be bounded open Lipschitz subsets in $\mathbf{R}^n$ such that $\Omega \subset \subset D_1 \subset \subset D$. Extend the function $\mathbf{u}$ to $\mathbf{R}^n$ by setting it to be zero outside Ω . From the argument in the proof of Theorem 4.5, it is clear that $\mathbf{u} \in D_0^1(D)$. Let $\tilde{\Omega} \equiv D_1 \setminus \overline{\Omega}$. Then $\tilde{\Omega} \subset \subset D$ is of class $\mathcal{C}$ since every bounded open Lipschitz set is of class $\mathcal{C}$, and $\mathbf{u} = 0$ over $\tilde{\Omega}$. Let $\tilde{\Omega}'$ be given as in Lemma 4.2 (i.e. $\tilde{\Omega}'$ can be retracted to $\tilde{\Omega}$) such that $\tilde{\Omega}' \subset \subset D$, and Let $\{V_j\}_{j=1}^m$ be the open covering of $\partial\Omega$, given as in (2.5) and (2.6), such that $\partial\Omega \subset \bigcup_{j=1}^m V_j \subset \subset \tilde{\Omega}'$. Then by Theorem 4.1' and Theorem 4.4', there are functions $\{\mathbf{u}_j\}_{j=1}^m$ and a neighborhood $\hat{U}$ of $\partial\Omega$ in $\mathbf{R}^n$ with $\hat{U} \subset \subset \bigcup_{j=1}^m V_j$ such that

$$\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j \quad \text{over } \hat{U}$$

and for all $j = 1, \dots, m$, $\mathbf{u}_j \in D_0^1(V_j)$, $\mathbf{u}_j = 0$ over $\tilde{\Omega}$. From which we imply $\mathbf{u}_j = 0$ a.e. in $D \setminus \overline{\Omega}$ for all $j = 1, \dots, m$.

Let $\mathbf{u}_0 = \mathbf{u} - \sum_{j=1}^m \mathbf{u}_j$. Then $\mathbf{u}_0 \in D_0^1(D)$ and $\mathbf{u}_0 = 0$ a.e. in $(D \setminus \overline{\Omega}) \cup \hat{U}$, from which we get $\mathbf{u}_0 = 0$ a.e. in $D \setminus (\Omega \setminus \overline{\hat{U}})$. Since $\Omega \setminus \overline{\hat{U}} \subset \subset \Omega$, one can find a Lipschitz open subset $\hat{\Omega}$ such that $\Omega \setminus \overline{\hat{U}} \subset \subset \hat{\Omega} \subset \subset \Omega$. Because $\mathbf{u}_0 = 0$ a.e. in $D \setminus \overline{\hat{\Omega}}$, we get by the trace theorem that $\mathbf{u}_0 \in D_0^1(\hat{\Omega})$. Then by Lemma 1.2.1 in Chapter 3 in [11], it follows that $\mathbf{u}_0 \in H_{0,\sigma}^1(\hat{\Omega}) \subset H_{0,\sigma}^1(\Omega)$.

Next for each $j = 1, \dots, m$, by setting $\mathbf{u}_j^t(x) = \mathbf{u}_j(x + t\mathbf{v}_j)$ for $t > 0$ small enough and using the same argument as that in the proof of Theorem 4.5, we obtain that $\mathbf{u}_j \in H_{0,\sigma}^1(\Omega)$.

Hence, we proved that $\mathbf{u} = \sum_{j=0}^m \mathbf{u}_j \in H_{0,\sigma}^1(\Omega)$. This completes the proof of Theorem 4.6. ■

Remark 4.7 After some slight modification in the proof of Theorem 3.1, we can verify a similar decomposition result for vector-valued function $\mathbf{u} \in L^2(D)^n$ with $\operatorname{div} \mathbf{u} = 0$ in D . (Here $\operatorname{div} \mathbf{u}$ is in the sense of distribution.) To be more precise, we let $\{U_j\}_{j=1}^m \subset D$ be open subsets such that $\bigcup_{j=1}^m U_j \subset \subset D$. Then for any open subset $\hat{U}$ with $\hat{U} \subset \subset \bigcup_{j=1}^m U_j$ there are functions $\{\mathbf{u}_j\}_{j=1}^m \subset L^2(D)^n$ such that $\mathbf{u} = \sum_{j=1}^m \mathbf{u}_j$ over $\hat{U}$, and for all $j = 1, \dots, m$, $\operatorname{div} \mathbf{u}_j = 0$ in D in the sense of distribution, $\operatorname{spt} \mathbf{u}_j \subset U_j$ and $\|\mathbf{u}_j\|_{L^2(D)} \leq C \|\mathbf{u}\|_{L^2(D)}$.

Furthermore, the same localization property as stated in Theorem 4.1 and Theorem 4.2 for such a partition still holds. Thus we can also obtain the following: If Ω is of class $\mathcal{C}$ in $\mathbf{R}^n$, $n = 2, 3$, then

$$L_\sigma^2(\Omega) = \{\mathbf{u} \in L^2(\mathbf{R}^n)^n; \operatorname{div} \mathbf{u} = 0 \text{ in } \mathbf{R}^n, \mathbf{u} = 0 \text{ a.e. in } \mathbf{R}^n \setminus \Omega\},$$

where $L_\sigma^2(\Omega) = \overline{C_{0,\sigma}^\infty(\Omega)}^{\|\cdot\|_{L^2}}$, the completion of $C_{0,\sigma}^\infty(\Omega)$ in the norm of $L^2(\Omega)^n$.

5. The existence of a shape optimization problem for the stationary Navier-Stokes equation

Let D and B be open bounded Lipschitz subsets in $\mathbf{R}^n$, $n = 2, 3$, such that $B \subset \subset D$. Let $\mathcal{O}$ be the family of certain open subsets Ω contained in B , which is defined as in section 2. Consider for each $\Omega \in \mathcal{O}$ the stationary Navier-Stokes equation on $D \setminus \overline{\Omega}$:

$$\begin{cases} -\gamma \Delta \mathbf{u} - (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p = \mathbf{f} & \text{in } D \setminus \overline{\Omega}, \\ \operatorname{div} \mathbf{u} = 0 & \text{in } D \setminus \overline{\Omega}, \\ u = 0 & \text{on } \partial D \cup \partial \Omega, \end{cases} \quad (5.1)$$

where $\mathbf{f}$ is a given function in $L^2(D)^n$. We shall denote by $\mathbf{u}_\Omega$ the weak solution of (5.1), i.e., $\mathbf{u}_\Omega \in H_{0,\sigma}^1(D \setminus \overline{\Omega})$ and

$$\gamma \int_{D \setminus \overline{\Omega}} \nabla \mathbf{u}_\Omega \cdot \nabla \Phi dx + \int_{D \setminus \overline{\Omega}} (\mathbf{u}_\Omega \cdot \nabla) \mathbf{u}_\Omega \cdot \Phi dx = \int_{D \setminus \overline{\Omega}} \mathbf{f} \cdot \Phi dx$$

for all $\Phi \in C_{0,\sigma}^\infty(D \setminus \overline{\Omega}) \equiv \{\Psi \in C_0^\infty(D \setminus \overline{\Omega})^n; \operatorname{div} \Psi = 0\}$.

It is well known that (see [11] and [6]) for each $\Omega \in \mathcal{O}$, (5.1) has at least one weak solution. Moreover, there is a positive constant $C(\gamma, d)$ depending only on γ and the width of D such that if

$$\|\mathbf{f}\|_{L^2(D)^n} < C(\gamma, d), \quad (5.2)$$

then the weak solution to (5.1) is unique. In the following we always assume that (5.2) holds.

Our purpose in this section is to study the following shape optimization problem:

$$(P) \quad \operatorname{Min}_{\Omega \in \mathcal{O}} \int_{B \setminus \overline{\Omega}} J(x, \mathbf{u}_\Omega, \nabla \mathbf{u}_\Omega) dx$$

subject to Equation (5.1).

Here we assume that J satisfies the following:

(H) $J : D \times \mathbf{R}^n \times \mathbf{R}^{n \times n} \rightarrow \mathbf{R}^+ \equiv [0, \infty)$ is measurable in the first variable and continuous in others. Moreover, there exist a positive constant C and a function $g \in L^1(D)$ such that for all $(x, \xi, \eta) \in D \times \mathbf{R}^n \times \mathbf{R}^{n \times n}$,

$$J(x, \xi, \eta) \leq C(g(x) + |\xi|_{\mathbf{R}^n}^2 + |\eta|_{\mathbf{R}^{n \times n}}^2). \quad (5.3)$$

Problem (P) is to ask the shape of a body Ω in $\mathcal{O}$ such that the cost functional takes its minimum. For instance, if we take $J(x, \mathbf{u}_\Omega, \nabla \mathbf{u}_\Omega) \equiv |\nabla \mathbf{u}_\Omega|^2 + |(\nabla \mathbf{u}_\Omega)^T|^2$, where $(\nabla \mathbf{u})^T$ denotes the transpose of $\nabla \mathbf{u}_\Omega$, then problem (P) is to ask the shape of a body Ω among $\mathcal{O}$ having the smallest drag. (see [10] .)

Theorem 5.1 Assume that the conditions in (4.2) and (4.3) are fulfilled. Then the shape optimization problem (P) has at least one solution $\Omega^* \in \mathcal{O}$.

Proof of Theorem 5.1 Let $\{\Omega_m\}_{m=1}^\infty \subset \mathcal{O}$ be a minimizing sequence for Problem (P). By Lemma 2.1, there exist some $\Omega^* \in \mathcal{O}$ and a subsequence of $\{\Omega_m\}_{m=1}^\infty$, still denoted by itself, such that $\operatorname{Hlim}_{m \rightarrow \infty} \Omega_m = \Omega^*$. We notice that $\Omega^* \subset \subset D$, for $B \subset \subset D$. Let $\mathbf{u}_m, m = 1, 2, \dots$, be the weak solutions to equation (4.1) with $\Omega = \Omega_m$. Then $\mathbf{u}_m \in H_{0,\sigma}^1(D \setminus \overline{\Omega}_m)$ and

$$\gamma \int_{D \setminus \overline{\Omega}_m} \nabla \mathbf{u}_m \cdot \nabla \Phi dx + \int_{D \setminus \overline{\Omega}_m} (\mathbf{u}_m \cdot \nabla) \mathbf{u}_m \cdot \Phi dx = \int_{D \setminus \overline{\Omega}_m} \mathbf{f} \cdot \Phi dx$$

for all $\Phi \in H_{0,\sigma}^1(D \setminus \overline{\Omega}_m)$. This immediately gives that

$$\int_{D \setminus \overline{\Omega}_m} |\nabla \mathbf{u}_m|^2 dx = \int_{D \setminus \overline{\Omega}_m} \mathbf{f} \cdot \mathbf{u}_m dx,$$

for $\int_{D \setminus \overline{\Omega}_m} (\mathbf{u}_m \cdot \nabla) \mathbf{u}_m \cdot \mathbf{u}_m dx = 0$. Thus

$$\int_{D \setminus \overline{\Omega}_m} |\nabla \mathbf{u}_m|^2 dx \leq C,$$

where C is a positive constant independent of m .

Let

$$\hat{\mathbf{u}}_m(x) = \begin{cases} \mathbf{u}_m(x) & \text{in } D \setminus \overline{\Omega}_m, \\ 0 & \text{in } \Omega_m \cup (\mathbf{R}^n \setminus D). \end{cases}$$

Then it is clear that $\hat{\mathbf{u}}_m \in H_{0,\sigma}^1(D \setminus \overline{\Omega}_m) \subset H_{0,\sigma}^1(D)$ and $\{\hat{\mathbf{u}}_m\}_{m=1}^\infty$ is bounded in $H_0^1(D)^n$. Thus there exists a subsequence of $\{\hat{\mathbf{u}}_m\}_{m=1}^\infty$, still denoted by itself, such that

$$\hat{\mathbf{u}}_m \rightarrow \hat{\mathbf{u}} \text{ weakly in } H^1(D)^n \text{ and strongly in } L^2(D)^n \quad (5.4)$$

for some $\hat{\mathbf{u}} \in H_{0,\sigma}^1(D)$.

We claim that $\hat{\mathbf{u}} = 0$ a.e. in Ω^* . Indeed, for any open subset K with $K \subset \subset \Omega^*$, it follows from Lemma 2.2 that there exists an integer $m_K > 0$ such that $\overline{K} \subset \Omega_m$ for all $m \leq m_K$. Thus

$$\int_K |\hat{\mathbf{u}}(x)|^2 dx = \lim_{m \rightarrow \infty} \int_K |\hat{\mathbf{u}}_m(x)|^2 dx \leq \overline{\lim}_{m \rightarrow \infty} \int_{\Omega_m} |\hat{\mathbf{u}}_m(x)|^2 dx = 0,$$

which implies $\hat{\mathbf{u}}(x) = 0$ a.e. in K and so $\hat{\mathbf{u}} = 0$ a.e. in Ω^* .

Now it follows from Theorem 4.5 that $\hat{\mathbf{u}} \in H_{0,\sigma}^1(D \setminus \overline{\Omega}^*)$.

We next prove

$$\gamma \int_{D \setminus \overline{\Omega}^*} \nabla \hat{\mathbf{u}} \cdot \nabla \Phi dx + \int_{D \setminus \overline{\Omega}^*} (\hat{\mathbf{u}} \cdot \nabla) \hat{\mathbf{u}} \cdot \Phi dx = \int_{D \setminus \overline{\Omega}^*} \mathbf{f} \cdot \Phi dx \quad (5.5)$$

for any $\Phi \in C_{0,\sigma}^\infty(D \setminus \overline{\Omega}^*)$. To this aim, we fix $\Phi \in C_{0,\sigma}^\infty(D \setminus \overline{\Omega}^*)$ and let $K = (spt \Phi)^o$, the interior of $spt \Phi$. Then by Lemma 2.3, there is a positive integer $m(K)$ such that $\overline{K} \subset D \setminus \Omega_m$ for any $m \geq m(K)$. Therefore,

$$\gamma \int_{spt \Phi} \nabla \hat{\mathbf{u}}_m \cdot \nabla \Phi dx + \int_{spt \Phi} \hat{\mathbf{u}}_m \cdot \Phi dx = \int_{spt \Phi} \mathbf{f} \cdot \Phi dx$$

for any $m \geq m(K)$. Passing to the limit, as $m \rightarrow \infty$, in the above, we get (5.5). Hence, $\hat{\mathbf{u}}$ is a weak solution to equation (5.1) where $\Omega = \Omega^*$.

Finally, we shall show

$$\int_{D \setminus \overline{\Omega}^*} J(x, \hat{\mathbf{u}}, \nabla \hat{\mathbf{u}}) dx = \text{Min}_{\Omega \in \mathcal{O}} \int_{D \setminus \overline{\Omega}} J(x, \mathbf{u}_\Omega, \nabla \mathbf{u}_\Omega) dx.$$

For this purpose, we let $D \setminus \overline{\Omega}^* = \bigcup_{j=1}^{\infty} G_j$ where $G_j, j = 1, \dots$, are open subsets in $D \setminus \overline{\Omega}^*$ such that $\overline{G_j} \subset G_{j+1}$. Then by Lemma 2.3, we obtain that for each j , there exists an integer $m_j > 0$ such that $\overline{G_j} \subset D \setminus \overline{\Omega}_m$ for any $m \geq m_j$. Therefore, we have for each j , that

$$\underline{\lim}_{m \rightarrow \infty} \int_{D \setminus \overline{\Omega}_m} J(x, \hat{\mathbf{u}}_m, \nabla \hat{\mathbf{u}}_m) dx \geq \underline{\lim}_{m \rightarrow \infty} \int_{G_j} J(x, \hat{\mathbf{u}}_m, \nabla \hat{\mathbf{u}}_m) dx. \quad (5.6)$$

On the other hand, since $\hat{\mathbf{u}}$ is the weak solution of (5.1) where $\Omega = \Omega^*$, we have

$$\gamma \int_{D \setminus \overline{\Omega}^*} |\nabla \hat{\mathbf{u}}|^2 dx = \int_{D \setminus \overline{\Omega}^*} \mathbf{f} \cdot \hat{\mathbf{u}} dx,$$

from which it follows that

$$\begin{aligned} \int_D |\nabla \hat{\mathbf{u}}_m|^2 dx - \int_D |\nabla \hat{\mathbf{u}}|^2 dx &= \int_{D \setminus \overline{\Omega}_m} |\nabla \mathbf{u}_m|^2 dx - \int_{D \setminus \overline{\Omega}^*} |\nabla \hat{\mathbf{u}}|^2 dx \\ &= \frac{1}{\gamma} \left(\int_{D \setminus \overline{\Omega}_m} \mathbf{f} \cdot \mathbf{u}_m dx - \int_{D \setminus \overline{\Omega}^*} \mathbf{f} \cdot \hat{\mathbf{u}} dx \right) \\ &= \frac{1}{\gamma} \int_D \mathbf{f} \cdot \hat{\mathbf{u}}_m dx - \int_D \mathbf{f} \cdot \hat{\mathbf{u}} dx \rightarrow 0 \quad \text{as } m \rightarrow \infty, \end{aligned}$$

i.e.,

$$\|\hat{\mathbf{u}}_m\|_{H^1(D)^n} \rightarrow \|\hat{\mathbf{u}}\|_{H^1(D)^n}.$$

This together with (5.4) implies that

$$\hat{\mathbf{u}}_m \rightarrow \hat{\mathbf{u}} \quad \text{strongly in } H^1(D)^n.$$

Now by the assumption in (H), Lemma 2.3 and by Fatou's lemma, we get for each j , the following:

$$\underline{\lim}_{m \rightarrow \infty} \int_{G_j} J(x, \hat{\mathbf{u}}_m, \nabla \hat{\mathbf{u}}_m) dx \geq \int_{G_j} J(x, \hat{\mathbf{u}}, \nabla \hat{\mathbf{u}}) dx.$$

This together with (5.6) shows that for each j fixed,

$$\begin{aligned} \text{Min}_{\Omega \in \mathcal{O}} \int_{D \setminus \overline{\Omega}} J(x, \mathbf{u}_\Omega, \nabla \mathbf{u}_\Omega) dx &\geq \underline{\lim}_{m \rightarrow \infty} \int_{G_j} J(x, \hat{\mathbf{u}}_m, \nabla \hat{\mathbf{u}}_m) dx \\ &\geq \int_{G_j} J(x, \hat{\mathbf{u}}, \nabla \hat{\mathbf{u}}) dx. \end{aligned}$$

Using Fatou's lemma again, we get

$$\begin{aligned} \text{Min}_{\Omega \in \mathcal{O}} \int_{D \setminus \overline{\Omega}} J(x, \mathbf{u}_\Omega, \nabla \mathbf{u}_\Omega) dx &\geq \underline{\lim}_{j \rightarrow \infty} \int_{D \setminus \overline{\Omega}^*} \chi_{G_j} J(x, \hat{\mathbf{u}}, \nabla \hat{\mathbf{u}}) dx \\ &\geq \int_{D \setminus \overline{\Omega}^*} \underline{\lim}_{j \rightarrow \infty} \chi_{G_j} J(x, \hat{\mathbf{u}}, \nabla \hat{\mathbf{u}}) dx \\ &= \int_{D \setminus \overline{\Omega}^*} J(x, \hat{\mathbf{u}}, \nabla \hat{\mathbf{u}}) dx. \end{aligned}$$

where χ_{G_j} denotes the characteristic function of G_j . Hence $\hat{\mathbf{u}}$ is an optimization solution for problem (P).

This completes the proof of Theorem 5.1. ■

Remark 5.2 By the same argument as above and making use of Theorem 4.6 instead, we can similarly obtain the existence for the following shape optimization problem:

$$(\tilde{P}) \quad \text{Min}_{\Omega \in \mathcal{O}} \int_{\Omega} J(x, \mathbf{u}_\Omega, \nabla \mathbf{u}_\Omega) dx$$

subject to

$$\begin{cases} -\gamma \Delta \mathbf{u} - (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p = \mathbf{f} & \text{in } \Omega, \\ \text{div} \mathbf{u} = 0 & \text{in } \Omega, \\ \mathbf{u} = 0 & \text{on } \partial\Omega, \end{cases} \quad (5.7)$$

where $\mathbf{u}_\Omega$ denotes the weak solution of (5.7) corresponding to $\Omega \in \mathcal{O}$.

References

- [1] R. A. Adams, *Sobolev Spaces*, Academic Press, New York, 1975.
- [2] R. Bott and Loring W. Tu, *Differential Forms in Algebraic Topology*, Springer-Verlag, New York, 1982.
- [3] D. Bucur and JP Zolesio, *N-dimensional shape optimization under capacity constraints*, J. Differential Equations, 123: 504-522, 1995.

- [4] D. Chenais, *On the existence of a solution in a domain identification problem*, J. Math. Anal. Appl., 52: 189-219, 1975.
- [5] Georges de Rham, *Differentiable Manifolds*, Translated from the French by F. R. Smith, Springer-Verlag, Heidelberg, 1984.
- [6] Giovanni P. Galdi, *An Introduction to The Mathematical Theory of The Navier-stokes Equations*, Springer-Verlag, New York, 1994.
- [7] L. Hedberg, *Spectral synthesis in Sobolev spaces and uniqueness of solution of the Dirichlet problem*, Acta Math. 147: 237-264, 1981.
- [8] V. Maz'ya, *Sobolev Spaces*, Springer-Verlag, Berlin, 1985.
- [9] Marvin J. Greenberg and John R. Harper, *Algebraic Topology, A First-Course*, The Benjamin/Cummings Publishing Company, London, 1981.
- [10] O. Pironneau, *Optimal Shape Design for Elliptic Systems*, Springer-Verlag, Berlin, 1984.
- [11] Hermann Sohr, *The Navier-Stokes Equations: An Elementary Functional Analytic Approach*, Birkhauser, Berlin, 2001.
- [12] V. Sverak, *On optimal shape design*, J. Math. Pures Appl., 72: 537-551, 1993.
- [13] R. Schneider, *Convex Bodies: The Brunn-Minkowski Theory*, Cambridge University Press, New York, 1993.
- [14] Dan Tiba, *A property of sobolev spaces and existence in optimal design*, Appl. Math. Optim., 47: 45-58, 2003.
- [15] G. S. Wang, L. Wang and D. H. Yang, *Shape optimization of elliptic equations in exterior domains*, submitted.